

RAMIFICATIONS, REDUCTION OF SINGULARITIES, SEPARATRICES

P. FORTUNY AYUSO

ABSTRACT. We study the behaviour of sequences of blowing-ups under ramifications, and use the results to give a simple proof of Camacho-Sad's Theorem on the existence of Separatrices for singularities of plane holomorphic foliations. The main result we prove is that for any finite sequence π of blowing-ups, there is a ramification morphism ρ such that the elimination of indeterminations $\tilde{\pi}$ of $\pi^{-1} \circ \rho$ is a sequence of blowing-ups *with centers at regular points of the exceptional divisors*; moreover, we show that if π is the reduction of singularities of a foliation $\mathcal{F}$, then ρ can be such that $\tilde{\pi}^* \rho^* \mathcal{F}$ has only simple singularities.

1. INTRODUCTION

In 1982, C. Camacho and P. Sad [3] proved that any holomorphic foliation in $(\mathbb{C}^2, 0)$ has at least one invariant analytic curve passing through $(0, 0)$. This result was searched for, at least, since XIX Century (cf. [1] in which it was proved for what are now called *simple* singularities). Afterwards, the same Camacho generalized it to vector fields in singular surfaces whose dual resolution graph is a tree [2]. In 1993, J. Cano [5] gave a constructive proof, which he simplified in 1997 [6]. Camacho and Sad's requires a detailed (and rather cumbersome) study of the combinatorial behaviour of what has since been called the *Camacho-Sad index* of an invariant manifold. J. Cano's, on the other hand, although being much simpler, introduces a somewhat artificial condition on that index.

Our work relies on the —new, as far as the author knows— study of the behaviour of sequences of blowing-ups under ramification morphisms. These maps were used successfully in [8] regarding Thom's Gradient Conjecture (see [10]) and [7] for studying some properties of the polar curves of a foliation. In the present paper we show —roughly speaking— how a sequence of blowing-ups can be modified by a ramification to obtain a new sequence *without combinatorial terms*: in the same way as one can modify a (singular) plane curve by a ramification in order to get a union of regular curves, we “modify” a sequence of blowing-ups in order to get another one which is “regular”.

2. REDUCTION OF SINGULARITIES AND RAMIFICATIONS

Recall that a foliation $\mathcal{F}$ in a smooth analytic surface $\mathcal{X}$ can be identified, *locally*, with a germ of holomorphic 1-form ω , whose local expression $a(x, y)dx + b(x, y)dy$ is such that a and b are relatively prime in $\mathbb{C}\{x, y\}$. We will (abusing notation) speak indifferently of $\mathcal{F}$ and ω when there is no possibility of confusion. A *separatrix* of $\mathcal{F}$ is a (germ of) holomorphic curve $\gamma : (\mathbb{C}, 0) \rightarrow \mathcal{X}$ such that the pull-back $\gamma^* \omega$

Date: February 1, 2008.

Supported by the European Commission Marie Curie Fellowship Programme.

is null. A point P is a *simple singularity* of ω if there are coordinates at P such that the linear part of the vector field $-b(x, y)\partial/\partial x + a(x, y)\partial/\partial y$ has two different eigenvalues $\mu \neq \lambda \neq 0$ with $\mu/\lambda \notin \mathbb{Q}_{>0}$.

Definition 1. A ramification morphism is a map $\rho : (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0)$ such that there are local analytic coordinates (u, v) and (x, y) at $(0, 0) \in (\mathbb{C}^2, 0)$ in which

$$(x(u, v), y(u, v)) = \rho(u, v) = (u^r, v)$$

for some positive integer r , called the ramification order. These coordinates will be called adapted to ρ .

Notice that we allow different changes of coordinates in each side of the map. If $\mathcal{F}$ is a germ of foliation in $(\mathbb{C}^2, 0)$, and $\rho^*\mathcal{F}$ is its pull-back by ρ , then any separatrix of $\rho^*\mathcal{F}$ is mapped by ρ into a separatrix of $\mathcal{F}$ (although, obviously, this assignment needs not be bijective).

All the arguments will be simplified if we introduce some notation: let $\pi : \mathcal{X} \rightarrow (\mathbb{C}^2, 0)$ be a sequence of blowing-ups $\pi = \pi_k \circ \cdots \circ \pi_1$ whose respective exceptional lines are E_i . We say that π is a *regular tree of blowing-ups* if all the centers are regular points of the exceptional divisor. The sequence π is a *chain of blowing-ups* if the center P_{i+1} of π_{i+1} belongs to E_i . Finally, a regular tree which is a chain will be called a *string of blowing-ups*. If π is a regular tree, we say that an irreducible component E_i of the exceptional divisor E is a *son of* component E_j (and that E_j is the father of E_i) if $E_i \cap E_j \neq \emptyset$ and $i > j$.

Given π as above and a ramification morphism ρ , we say that ρ is *transversal to π* if there are coordinates (u, v) and (x, y) adapted to ρ such that none of the centers of π_i is the infinitely near point given by $T_{(0,0)}(x = 0)$.

Recall [9] that given a rational function $g/h = f : \mathcal{Y} \rightarrow \mathbb{C}$ where $\mathcal{Y}$ is an analytic surface, a point $P \in \mathcal{Y}$ is an *indetermination point of f* if $g(P) = h(P) = 0$. It is well-known that there is a finite sequence of point blowing-ups $\eta : \tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$ such that $f \circ \eta$ is well-defined everywhere; this η is the *elimination of indeterminations of f* . Obviously, this construction can be applied to maps from $\mathcal{Y}$ to any analytic space (see [9] for the details).

As our aim is to prove the Separatrix Theorem, we are going to assume from now on that all our foliations are *non-dicritical*.

Our main result is the following

Theorem 2. Let $\pi : \mathcal{X} \rightarrow (\mathbb{C}^2, 0)$ be the minimal reduction of singularities of $\mathcal{F}$. There exists a ramification ρ such that if $\tilde{\pi} : \tilde{\mathcal{X}} \rightarrow (\mathbb{C}^2, 0)$ is the elimination of indeterminations of $\pi^{-1} \circ \rho : (\mathbb{C}^2, 0) \rightarrow \mathcal{X}$, then $\tilde{\pi}$ is a regular tree and all the singularities of $\tilde{\pi}^*\rho^*\omega$ are simple.

As an obvious consequence, we get:

Corollary 3. There is a ramification morphism ρ such that the reduction of singularities of $\rho^*\omega$ is a regular tree.

Before giving the proof of Theorem 2, it is convenient to introduce some more notation:

Definition 4. Let $E = \cup_{i=1}^n E_i$ be the exceptional divisor of π . We say that the irreducible component E_i corresponding to π_i is *$\mathcal{F}$ -terminal* if there is no $j > i$ such that the center P_j of π_j belongs to E_i .

A germ of analytic curve $\gamma : (\mathbb{C}, 0) \rightarrow (\mathbb{C}^2, 0)$ has the *type of singularity* of E_i if its strict transform $\pi^*\gamma$ meets E transversely at E_i (as a consequence, $\pi^*\gamma$ is also regular). The germ γ is $\mathcal{F}$ -terminal if it has the type of singularity of an $\mathcal{F}$ -terminal divisor.

Remark 5. It is clear that the reduction of singularities of a foliation $\mathcal{F}$ is a regular tree if and only if any $\mathcal{F}$ -terminal curve is regular.

We shall make constant use of the following result, whose proof is an easy computation:

Lemma 6. *Let π be a sequence of blowing-ups and ρ a ramification of order r transversal to π . Call $\tilde{\pi}$ to the elimination of singularities of $\pi^{-1} \circ \rho$. Then:*

- a) *If π is a string of n blowing-ups then $\tilde{\pi}$ is a string of nr blowing-ups.*
- b) *As a consequence, if π is a tree with b branches of lengths $l_1, \dots, l_b$, then $\tilde{\pi}$ is a tree with b branches of lengths $rl_1, \dots, rl_b$.*

For the reader's sake, we give the local expression of $\tilde{\rho} = \pi^{-1} \circ \rho \circ \tilde{\pi}$ when π is a string. Fix systems of coordinates (u, v) and (x, y) in $(\mathbb{C}^2, 0)$ adapted to ρ . Cover $\mathcal{X}$ and $\tilde{\mathcal{X}}$ as follows:

$$(1) \quad \begin{cases} \mathcal{X}_n = (\overline{U}_1 \cup \dots \cup \overline{U}_n) \cup U_n \\ \tilde{\mathcal{X}} = (\overline{U}_{11} \cup \dots \cup \overline{U}_{1r}) \cup \dots \cup (\overline{U}_{n1} \cup \dots \cup \overline{U}_{nr}) \cup U_{nr} \end{cases}$$

where each $\overline{U}_i$ is the standard chart covering the point at infinity of the exceptional divisor E_i of π_i , and U_n covers the origin of E_n . The open set $\overline{U}_{ij}$ covers the point at infinity of $E_{r(i-1)+j}$ and U_{nr} covers the origin of E_{nr} . These charts have respective systems of coordinates $(\overline{x}_i, \overline{y}_i)$ and (x_n, y_n) for the $\overline{U}_i$'s and U_n , and $(\overline{u}_{ij}, \overline{v}_{ij})$ and (u_{nr}, v_{nr}) for the $\overline{U}_{ij}$ and U_{nr} , which can be chosen (in a standard way) so that

$$(2) \quad \begin{cases} (\overline{u}_{nr}, \overline{v}_{nr}) = \left(\frac{1}{v_{nr}}, u_{nr}v_{nr} \right) \text{ and} \\ (\overline{u}_{i'j'}, \overline{v}_{i'j'}) = \left(\frac{1}{v_{ij}}, u_{ij}v_{ij}^2 \right) \text{ when } r(i-1) + j = r(i'-1) + j' + 1 \end{cases}$$

and the analog conditions in $\mathcal{X}$. We have (assuming π is a string):

Proposition 7. *Let $\tilde{E} = (E_{11} \cup \dots \cup E_{1r}) \cup \dots \cup (E_{n1} \cup \dots \cup E_{nr})$ be the exceptional divisor of $\tilde{\pi}$ and $E = E_1 \cup \dots \cup E_n$ that of π . Call P_i to the origin of the chart $\overline{U}_i$ (the point "at infinity"). Then:*

- 1. *For $i = 1, \dots, n$ and $j = 1, \dots, r-1$, $\tilde{\rho}(E_{ij}) = P_i$.*
- 2. *For $i = 1, \dots, n$, $\tilde{\rho}(E_{ir}) = E_i$.*

Moreover, given i and j as above, The local expression of $\tilde{\rho}$ in $\overline{U}_{ij}$ and $\overline{U}_i$ is:

$$(3) \quad (\overline{x}_i, \overline{y}_i) = \tilde{\rho}(\overline{u}_{ij}, \overline{v}_{ij}) = \left(\overline{u}_{ij}^{r-j+1} \overline{v}_{ij}^{r-j}, \overline{u}_{ij}^{j-1} \overline{v}_{ij}^j \right)$$

Combining Proposition 7 with Lemma 6 and taking into account that the transform of a simple singularity by a map of the form (3) is also a simple singularity, one infers:

Corollary 8. *If ρ is such that the reduction of singularities of $\rho^*\mathcal{F}$ is a regular tree, then for any other ramification σ , the reduction of singularities of $(\rho \circ \sigma)^*\mathcal{F}$ is also a regular tree.*

Proof of Theorem 2. Let $T = F_1 \cup \dots \cup F_p$ be the union of all the terminal divisors of $\mathcal{F}$, and let $\gamma_1, \dots, \gamma_p$ be a corresponding family of $\mathcal{F}$ -terminal curves. We can assume, by Lemma 6 that all the γ_i are singular, and that π is given by the sequence $\pi = \pi_p \circ \dots \circ \pi_1$, where π_i is (the remaining part of) the sequence of blowing-ups leading to T_i . Again by Lemma 6 and Remark 5, it is enough to show that for any i , there is a ramification morphism ρ_i such that $\rho_i^* \gamma_i$ is a union of regular curves, for $\rho = \rho_p \circ \dots \circ \rho_1$ would satisfy the thesis. Hence, we need only prove that given an irreducible singular curve γ , there is a ramification ρ such that

1. The curve $\rho^* \gamma$ is a union of non-singular branches.
2. The elimination of indeterminations of $\pi_1^{-1} \circ \rho$, say $\tilde{\pi} : \tilde{\mathcal{X}} \rightarrow (\mathbb{C}^2, 0)$, is a regular tree.
3. If $Q \in \tilde{\mathcal{X}}$ is such that $\tilde{\rho}(Q)$ is simple for $\pi^* \omega$, then Q is simple for $\tilde{\rho}^* \pi_i^* \omega$.

If γ has multiplicity m , then property (1) holds obviously for any ramification of order r multiple of m .

Let Γ be a curve with only one Puiseux exponent, having maximal contact with γ , so that they share all the infinitely near points of the reduction of singularities π_Γ of Γ . Let T_Γ be the last exceptional divisor of π_Γ . Reasoning by induction on the number of Puiseux exponents of γ and using Lemma 6 again, we may assume that γ has a single Puiseux exponent, m/n with $(m, n) = 1$.

Let then ρ be a ramification morphism of order r transversal to γ and write the continuous fraction expansion of $m/n = [a_0, a_1, \dots, a_s]$. Let π_0 be the string following the maximal contact sequence of γ , π^0 the remaining blowing-ups and consider the diagram

$$(4) \quad \begin{array}{ccc} & & \mathcal{X} \\ & \nearrow \tilde{\rho} & \downarrow \pi^0 \\ \tilde{\mathcal{X}} & \xrightarrow{\tilde{\rho}_0} & \mathcal{X}_0 \\ \downarrow \tilde{\pi}_0 & & \downarrow \pi_0 \\ B_2 & \xrightarrow{\rho} & B_2 \end{array}$$

where $\tilde{\mathcal{X}}$ is the elimination of singularities of $\pi_0^{-1} \circ \rho$. We use the notation of Proposition 7 for π_0 . We want to prove that for some r , the map $\tilde{\rho}_0$ lifts holomorphically to $\mathcal{X}$ (in other words, the *rational* map $\tilde{\rho}$ is actually holomorphic). The sequence π^0 starts by blowing-up $E_n \cap E_{n-1}$, so that we need only check that $\tilde{\rho}_0$ *restricted* to $E_{n-1r} \cup E_{n1} \cup \dots \cup E_{nr-1}$ lifts to $\mathcal{X}$. Consider the matrices

$$A = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad B = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix},$$

and, for any $i = a_1 + \dots + a_l + s$ let $\varphi_i(A, B) = B^s A^{a_l} \dots B^{a_2} A^{a_1}$ (assuming l odd, and the corresponding product for l even). The equations of $\tilde{\rho}$ at U_{n_j} are

$$(5) \quad \tilde{\rho}(u_j, v_j) = (u_j^{e_1} v_j^{e_2}, u_j^{f_1} v_j^{f_2})$$

where

$$\begin{pmatrix} e_1 & e_2 \\ f_1 & f_2 \end{pmatrix} = \varphi_i(A, B) \cdot \begin{pmatrix} r-j+1 & r-j \\ j-1 & j \end{pmatrix}$$

for some $0 \leq i \leq a_1 + \dots + a_s$ which depends on the point of $\mathcal{X}$ we are looking at. From this, one sees that $\tilde{\rho}$ is holomorphic in U_{n_j} if and only if $e_1 e_2 \geq 0$ and

$f_1 f_2 \geq 0$ (i.e. if the exponents in each component do not have different signs). We are seeking for an r for which all the possible pairs (e_1, e_2) and (f_1, f_2) share this property. Another inductive argument shows that for all i , there exist integer numbers c_i, d_i, m_i such that

$$\begin{pmatrix} e_1 & e_2 \\ f_1 & f_2 \end{pmatrix} = \begin{pmatrix} r - c_i(m_i - 1) & r - c_i m_i \\ d_i(m_i - 1) & d_i m_i \end{pmatrix}.$$

Hence, if r is a multiple of $c_i d_i$ for all $0 \leq i \leq a_1 + \dots + a_s$, then all the above maps are well-defined and $\tilde{\rho}$ is holomorphic. As the number of i 's is finite, this r exists. $\square$

3. RECALLING RESIDUES

Given a germ of holomorphic foliation $\mathcal{F}$ in $(\mathbb{C}^2, 0)$, and a regular separatrix E passing through $Q = (0, 0)$, the *Camacho-Sad* index of $\mathcal{F}$ at Q along E is defined as

$$I_Q(\mathcal{F}, S) = -\operatorname{Res}_0 \left(\frac{a(x, 0)}{b(x, 0)} dx \right).$$

where we are assuming that $\mathcal{F}$ is defined near Q by $\omega = ya(x, y)dx + b(x, y)dy$ and $E \equiv (y = 0)$.

One key result for any (known) proof of the Separatrix Theorem is the following classical result (see [4] for a complete proof and references to the original sources):

Proposition 9. *If Q is a simple singularity of $\mathcal{F}$, S is a smooth separatrix of $\mathcal{F}$ through Q with $I_Q(\mathcal{F}, S) \neq 0$ then there is another (smooth) separatrix T passing through Q .*

Let now $\pi : \mathcal{X} \rightarrow (\mathbb{C}^2, 0)$ be the blowing-up of $(0, 0)$ and call E to the exceptional divisor. Then

Theorem 10 ([3]). *If E is invariant for the pull-back $\pi^*\mathcal{F}$ of $\mathcal{F}$ and $P_1, \dots, P_r$ are the singular points of $\pi^*\mathcal{F}$ in E , then*

$$(6) \quad \sum_{i=1}^r I_{P_i}(\mathcal{F}, S) = -1$$

Moreover, if S is invariant for $\mathcal{F}$, π^*S is its pull-back by π and $Q = E \cap S$, then

$$(7) \quad I_Q(\pi^*\mathcal{F}, \pi^*S) = I_{(0,0)}(\mathcal{F}, S) - 1.$$

Finally, it is easily verified that if S and T are both smooth separatrices of $\mathcal{F}$ and $Q = S \cap T$ is a simple singularity of $\mathcal{F}$, then

$$(8) \quad I_Q(\mathcal{F}, S) = \frac{1}{I_Q(\mathcal{F}, T)}$$

whenever any of both indices is *non-zero*, or using the standard terminology, when Q is not a saddle-node singularity of $\mathcal{F}$.

4. PROOF OF THE SEPARATRIX THEOREM

We have now all the machinery needed to give a straightforward proof of the Separatrix Theorem:

Theorem 11 ([3]). *Given a germ of holomorphic foliation $\mathcal{F}$ at $(\mathbb{C}^2, 0)$, there is a separatrix for $\mathcal{F}$ passing through $(0, 0)$.*

Proof. As stated in Section 2, ramifications send separatrices of $\rho^*\mathcal{F}$ into separatrices of $\mathcal{F}$ so that, by Corollary 3, we need only prove the result for foliations whose reduction of singularities is a regular tree.

Assume then, that the reduction of singularities $\pi : \mathcal{X} \rightarrow (\mathbb{C}^2, 0)$ of $\mathcal{F}$ is a tree and let $E = E_1 \cup \dots \cup E_n$ be the exceptional divisor. By Proposition 9, we only need to show that there is a point $Q \in E_i$ for some i with $Q \notin E_j$ for $i \neq j$ such that $I_Q(\mathcal{F}, E_i) \neq 0$.

Assume, by contradiction, that there is no such Q . That is, for any Q in the regular part of E , the corresponding index is 0. We have: **Assertion:** Given an irreducible component F of E , if $F_1, \dots, F_s$ are its sons (s may be 0), then

$$(9) \quad \sum_{Q \in F} I_Q(\mathcal{F}, F) = -s - 1,$$

and all the terms in the sum are *rational numbers*. This is (easily) proved by induction on the maximal length of a branch starting from F , using Theorem 10 and our assumption that all the residues at the regular part of F are 0. A direct consequence of this is that if F' is the father of F and $P = F \cap F'$, then

$$(10) \quad I_Q(\mathcal{F}, F') \geq -1$$

(and the above index is a rational number, in fact).

But now, the divisor E_1 appearing in the first blowing-up has no father, so that all the singularities belonging to it either are the crossing with a son or have zero index along E_1 . From the Assertion and Equation (10), if $F_1, \dots, F_s$ are the sons of E_1 and $Q_i = E_1 \cap F_i$, we must have

$$\begin{aligned} -r - 1 &\stackrel{(9)}{=} \sum I_Q(\mathcal{F}, E_1) \\ &= \sum_{i=1}^s I_{Q_i}(\mathcal{F}, E_1) \\ &\stackrel{(10)}{\geq} \sum_{i=1}^s -1 = -s \end{aligned}$$

which gives the desired contradiction. $\square$

REFERENCES

- [1] C. A. Briot and J. C. Bouquet, *Propriétés des fonctions définies par des equations différentielles*, Journal de L'École Polytechnique (1856), no. 36, 133–198.
- [2] C. Camacho, *Quadratic forms and holomorphic foliations on singular surfaces*, Math. Ann. (1988), no. 282, 177–184.
- [3] C. Camacho and P. Sad, *Invariant varieties through singularities of holomorphic vector fields*, Ann. of Math. **115** (1982), 579–595.
- [4] C. Camacho and P. Sad, *Pontos singulares de equações diferenciais analíticas*, 16 Colóquio Brasileiro de Matemática, IMPA, Rio de Janeiro, 1985.
- [5] J. Cano, *An extension of the Newton-Puiseux polygon construction to give solutions of pfaffian forms*, Ann. de L'Institut Fourier (1993), no. 43, 125–142.
- [6] ———, *Construction of invariant curves for singular holomorphic vector fields*, Proc. of the AMS **125** (1997), no. 9, 2649–2650.
- [7] N. Corral, *Polar curves of singular foliations*, Ph.D. thesis, Univ. de Valladolid, 2001.
- [8] P. Fortuny and F. Sanz, *Gradient vector fields do not generate twister dynamics*, Journal of Differential Equations **174** (2001), 91–100.
- [9] H. Hironaka, *Resolution of singularities of an algebraic variety over a field of characteristic zero*, Ann. of Math. **79** (1964), 109–306.

- [10] K. Kurdyka, T. Mostowski, and A. Parusinski, *Proof of the Gradient Conjecture of R. Thom*, To appear in *Ann. of Math.*

DEPARTMENT OF MATHEMATICAL SCIENCES. SCHOOL OF MATHEMATICS. QUEEN MARY COLLEGE, UNIVERSITY OF LONDON. MILE END ROAD. E1 4NS LONDON, UK.

E-mail address: `p.fortuny@maths.qmul.ac.uk`